

Request for Prior Authorization JANUS KINASE (JAK) INHIBITORS

FAX Completed Form To 1 (800) 574-2515 Provider Help Desk 1 (877) 776-1567

(PLEASE PRINT - ACCURACY IS IMPORTANT)

IA Medicaid Member ID #	Patient name	DOB
Patient address		
Provider NPI	Prescriber name	Phone
Prescriber address		Fax
Pharmacy name	Address	Phone
-		
Prescriber must complete all information	tion above. It must be legible, correct, and complete or	form will be returned.
Pharmacy NPI	Pharmacy fax NDC	

Prior authorization (PA) is required for Janus kinase (JAK) inhibitors. Requests for non-preferred agents may be considered when documented evidence is provided that the use of the preferred agent(s) would be medically contraindicated. Payment will be considered for an FDA approved or compendia indicated diagnosis for the requested drug, excluding requests for the FDA approved indication of alopecia areata or other excluded medical use(s), as defined in Section 1927(d)(2) of the Social Security Act, State Plan, and Rules when the following conditions are met:

- I. Patient is not using or planning to use a JAK inhibitor in combination with other JAK inhibitors, biological therapies, or potent immunosuppressants (azathioprine or cyclosporine); and
- 2. Request adheres to all FDA approved labeling for requested drug and indication, including age, dosing, contraindications, warnings and precautions, drug interactions, and use in specific populations; and
- 3. Patient has a diagnosis of:
 - a. Moderate to severe rheumatoid arthritis (baricitinib, tofacitinib, upadacitinib); with
 - i. A documented trial and inadequate response, at a maximally tolerated dose, with methotrexate; and
 - ii. A documented trial and inadequate response to one preferred TNF inhibitor; or
 - b. Psoriatic arthritis (tofacitinib, upadacitinib); with
 - i. A documented trial and inadequate response, at a maximally tolerated dose, with methotrexate (leflunomide or sulfasalazine may be used if methotrexate is contraindicated); and
 - ii. Documented trial and therapy failure with one preferred TNF inhibitor used for psoriatic arthritis; or
 - c. Moderately to severely active ulcerative colitis (tofacitinib, upadacitinib); with
 - i. A documented trial and inadequate response with a preferred TNF inhibitor; and
 - ii. If requested dose for tofacitinib is 10mg twice daily, an initial 16 weeks of therapy will be allowed. Continued requests as this dose will need to document an adequate therapeutic benefit; or
 - d. Moderately to severely active Crohn's disease (upadacitinib); with
 - i. A documented trial and inadequate response with a preferred TNF inhibitor; or
 - e. Polyarticular Course Juvenile Idiopathic Arthritis (tofacitinib); with
 - i. A documented trial and inadequate response to the preferred oral DMARD, methotrexate (leflunomide or sulfasalazine may be used if methotrexate is contraindicated); and
 - ii. A documented trial and inadequate response with a preferred TNF inhibitor; or
 - f. Axial spondyloarthritis conditions (e.g., ankylosing spondylitis or nonradiographic axial spondyloarthritis) (tofacitinib, upadacitinib); with
 - i. A documented trial and inadequate response to at least two preferred non-steroidal anti-inflammatories (NSAIDS) at a maximally tolerated dose for a minimum of at least one month; and
 - ii. A documented trial and inadequate response with at least one preferred TNF inhibitor; or
 - g. Atopic dermatitis; with
 - i. Documentation patient has failed to respond to good skin care and regular use of emollients; and
 - ii. A documented adequate trial and therapy failure with one preferred medium to high potency topical corticosteroid for a minimum of 2 consecutive weeks; or
 - iii. A documented trial and therapy failure with a topical immunomodulator for a minimum of 4 weeks; and
 - iv. For mild to moderate atopic dermatitis (ruxolitinib):
 - a. Affected area is less than 20% of body surface area (BSA); and
 - b. Patient has been instructed to use no more than 60 grams of topical ruxolitinib per week; or
 - v. For moderate to severe atopic dermatitis (abrocitinib, upadacitinib):
 - a. A documented trial and therapy failure with a systemic drug product for the treatment of moderate to severe atopic dermatitis, including biologics; and
 - Requests for upadacitinib for pediatric patients 12 to less than 18 years if age must include the patient's weight in kg; or
 - h. Nonsegmental vitiligo (ruxolitinib) with;
 - i. A documented trial and inadequate response with a potent topical corticosteroid; or
 - ii. A documented trial and inadequate response with a topical calcineurin inhibitor; and
 - iii. The patient's body surface area (BSA) is less than or equal to the affected BSA per FDA approved label, if applicable.

The required trials may be overridden when documented evidence is provided that use of these agents would be medically contraindicated.

470-5175 (Rev. 1/25) Page 1 of 3

Request for Prior Authorization JANUS KINASE (JAK) INHIBITORS (PLEASE PRINT – ACCURACY IS IMPORTANT)

<u>Preferred</u>			Non-Preferr	<u>ed</u>		
□ Rinvoq □	Opzelura	□ Xeljanz	☐ Cibinqo	□ Olumiant	□ Xeljanz Or	ral Solution ☐ Xeljanz XR
Strength	Dos	age Instructions		Qu	antity	Days Supply
Diagnosis: _						
		e used in combin sants?		ther JAK inhil	bitors, biolog	ical therapies or
	to Severe R	heumatoid Arth	ritis (RA) (Ol	umiant, Rinvo	oq, Xeljanz or	Xeljanz XR)
						ates:
		ame/Dose:				ates:
☐ Psoriatic	Arthritis (Ri	nvoq, Xeljanz or	Xeljanz XR)			
		nide or sulfasalaz				.
						tes:
Preferred TNF	Inhibitor: N				Trial Da	ates:
		nvoq, Xeljanz or				
Preferred TNF	Inhibitor: N	ame/Dose:			Trial D	ates:
Failure reason	:					
		tofacitinib 10mg tw	•	·		c benefit:
	ly to severel	y active Crohn's	disease (Ri	nvoq)		
Preferred TNF Inhibitor: Name/Dose: Failure reason:					ates:	
☐ Polyarticu	ular Course	Juvenile Idiopat	hic Arthritis	(Xeljanz)		
		nide or sulfasalaz				dates:
Failure reason	:					
Preferred TNF	Inhibitor: N	ame/Dose:			Trial [Dates:
Failure reason	:					
		is conditions (e.; is) (Rinvoq, Xelja			or nonradio	graphic
Preferred NS	AID trial 1: Na	me/Dose:			_ Trial Dates:	
Fallons na a · · · ·	_					

Page 2 of 3 470-5175 (Rev. 1/25)

Request for Prior Authorization JANUS KINASE (JAK) INHIBITORS

(PLEASE PRINT - ACCURACY IS IMPORTANT)

	Trial dates:
	Trial Dates:
Failure reason:	
☐ Atopic Dermatitis Has patient failed to respond to good skin care and	d regular use of emollients? Yes No
Document emollient use: Product name, dosing instruc	ctions & duration of use:
Document trial and therapy failure with one preferred nweeks or topical immunomodulator for a minimum of 4	nedium to high potency topical corticosteroid for a minimum of 2 weeks:
Preferred Medium to High Potency Topical Cortico: Drug name & dose:	Trial dates:
Preferred Topical Immunomodulator Trial: Drug name & dose:	Trial dates:
Mild to Moderate Atopic Dermatitis (Opzelura)	
ls affected area less than 20% of body surface area	n? □Yes □No
Is affected area less than 20% of body surface area	
Has patient been instructed to use no more than 60	Ogms of topical ruxolitinib per week? Yes No
Has patient been instructed to use no more than 60 Moderate to Severe Atopic Dermatitis (Cibinqo or F	Ogms of topical ruxolitinib per week? Yes No
Has patient been instructed to use no more than 60 Moderate to Severe Atopic Dermatitis (Cibingo or F Trial with systemic drug product for the treatment of mo	Ogms of topical ruxolitinib per week? Yes No
Has patient been instructed to use no more than 60 Moderate to Severe Atopic Dermatitis (Cibinqo or F Trial with systemic drug product for the treatment of mo	Ogms of topical ruxolitinib per week? Yes No Rinvoq) oderate to severe atopic dermatitis, including biologics: Trial dates:
Has patient been instructed to use no more than 60 Moderate to Severe Atopic Dermatitis (Cibingo or F Trial with systemic drug product for the treatment of mo	Ogms of topical ruxolitinib per week? Yes No Rinvoq) oderate to severe atopic dermatitis, including biologics: Trial dates:
Has patient been instructed to use no more than 60 Moderate to Severe Atopic Dermatitis (Cibinqo or F Trial with systemic drug product for the treatment of mo Drug name & dose: Failure reason:	Ogms of topical ruxolitinib per week? Yes No Rinvoq) oderate to severe atopic dermatitis, including biologics: Trial dates:
Has patient been instructed to use no more than 60 Moderate to Severe Atopic Dermatitis (Cibinqo or Formaticial with systemic drug product for the treatment of moderate and the control of the treatment of moderate reason: Requests for upadacitinib for pediatric patients 12 Nonsegmental vitiligo (Opzelura) Potent Topical Corticosteroid Trial: Drug name & dose:	Ogms of topical ruxolitinib per week?
Has patient been instructed to use no more than 60 Moderate to Severe Atopic Dermatitis (Cibinqo or Formaticial with systemic drug product for the treatment of moderate and the control of the treatment of moderate reason: Requests for upadacitinib for pediatric patients 12 Nonsegmental vitiligo (Opzelura) Potent Topical Corticosteroid Trial: Drug name & dose:	Ogms of topical ruxolitinib per week?
Has patient been instructed to use no more than 60 Moderate to Severe Atopic Dermatitis (Cibinqo or F Trial with systemic drug product for the treatment of moderate to severe Atopic Dermatitis (Cibinqo or F Trial with systemic drug product for the treatment of moderate reason: Failure reason: Nonsegmental vitility (Opzelura) Potent Topical Corticosteroid Trial: Drug name & dose: Failure reason: Topical Calcineurin Inhibitor Trial:	Ogms of topical ruxolitinib per week?
Has patient been instructed to use no more than 60 Moderate to Severe Atopic Dermatitis (Cibinqo or F Trial with systemic drug product for the treatment of moderate to severe Atopic Dermatitis (Cibinqo or F Trial with systemic drug product for the treatment of moderate reason: Failure reason: Nonsegmental vitility (Opzelura) Potent Topical Corticosteroid Trial: Drug name & dose: Failure reason: Topical Calcineurin Inhibitor Trial:	Ogms of topical ruxolitinib per week?
Has patient been instructed to use no more than 60 Moderate to Severe Atopic Dermatitis (Cibinqo or Failure reason: Requests for upadacitinib for pediatric patients 12 Nonsegmental vitiligo (Opzelura) Potent Topical Corticosteroid Trial: Drug name & dose: Failure reason: Topical Calcineurin Inhibitor Trial: Drug name & dose: Failure reason:	Ogms of topical ruxolitinib per week?
Has patient been instructed to use no more than 60 Moderate to Severe Atopic Dermatitis (Cibinqo or Firial with systemic drug product for the treatment of moderate and a dose: Failure reason: Requests for upadacitinib for pediatric patients 12 Nonsegmental vitiligo (Opzelura) Potent Topical Corticosteroid Trial: Drug name & dose: Failure reason: Topical Calcineurin Inhibitor Trial: Drug name & dose: Failure reason: Provide patient's affected body surface area (BSA)	Ogms of topical ruxolitinib per week?

IMPORTANT NOTE: In evaluating requests for prior authorization the consultant will consider the treatment from the standpoint of medical necessity only. If approval of this request is granted, this does not indicate that the member continues to be eligible for Medicaid. It is the responsibility of the provider who initiates the request for prior authorization to establish by inspection of the member's Medicaid eligibility card and, if necessary by contact with the county Department of Health and Human Services, that the member continues to be eligible for Medicaid.

470-5175 (Rev. 1/25) Page 3 of 3